

References

1. Statistisches Bundesamt: Statistisches Jahrbuch 2014. URL <https://www.de-statistik.de/DE/Publikationen/Statistisches-Jahrbuch/StatistischesJahrbuch.html> – Zugriff:11.01.2015
2. Oberritter H, Schäbenthal K, von Ruesten A, et al. (2013) *The DGE-Nutrition Circle – Presentation and Basis of the Food-Related Recommendations from the German Nutrition Society (DGE). Ernährungs Umschau international.* 60 (2): 24–29
3. Bergleiter, S (2012) Nachhaltiger Fischkonsum: Ist die Empfehlung der DGE zum Fischverzehr unter Nachhaltigkeitsaspekten vertretbar? *Ernährungsumschau* 5:282–285
4. Souci SW, Fachmann W, Kraut H. *Die Zusammensetzung der Lebensmittel, Nährwert Tabellen.* 7. Auflage, MedPharm Scientific Publishers, Stuttgart (2008)
5. Galli C, Rise P (2009) Fish consumption, omega 3 fatty acids and cardiovascular disease. *The science and the clinical trials. Nutrition and health.* 20 (1): 11–20
6. König A, Bouzan C, Cohen JT, et al. (2005) A quantitative analysis of fish consumption and coronary heart disease mortality. *AM J PREV MED.* 29 (4): 335–346
7. *Noncommunicable diseases country profiles.* Geneva, Switzerland : World Health Organization. (2011)
8. Rapsomaniki E, Timmis A, George J, et al. (2014) Blood pressure and incidence of twelve cardiovascular diseases: lifetime risks, healthy life-years lost, and age-specific associations in 1.25 million people. *Lancet.* 383 (9932): 1899–1911
9. Dokholya RS, Albert CM, Appel LJ, et al. (2004) A trial of omega-3 fatty acids for prevention of hypertension. *AM J CARDIOL.* 93 (8): 1041–1043
10. Geleijnse JM, Giltay EJ, Grobbee DE, et al. (2002) Blood pressure response to fish oil supplementation: metaregression analysis of randomized trials. *J HYPERTENS.* 20 (8): 1493–1499
11. Mori TA, Burke V, Puddey IB, et al. (2009) The effects of omega 3 fatty acids and coenzyme Q(10) on blood pressure and heart rate in chronic kidney disease: a randomized controlled trial. *J HYPERTENS.* 27(9): 1863–1872
12. Ramel A, Alfredo MJ, Kiely M et al. (2010) Moderate consumption of fatty fish reduces diastolic blood pressure in overweight and obese European young adults during energy restriction. *Nutrition.* 26 (2):168–174
13. Theobald HE, Goodall AH, Sattar N, et al. (2007) Low-dose docosahexaenoic acid lowers diastolic blood pressure in middle-aged men and women. *J NUTR.* 137 (4): 973–978
14. Hoshi T, Wissuwa B, Tian Y, et al. (2013) Omega-3 fatty acids lower blood pressure by directly activating large-conductance Ca²⁺-dependent K(+)channels. *P NATL ACAD SCI USA.* 110 (12): 4816–4821
15. Liu JC, Conklin SM, Manuck SB, et al. (2011) Long-Chain Omega-3 Fatty Acids and Blood Pressure. *AM J HYPERTENS.* 24 (10): 1121–1126
16. Lara JJ, Economou M, Wallace AM, et al. (2007) Benefits of salmon eating on traditional and novel vascular risk factors in young, non-obese healthy subjects. *ATHEROSCLEROSIS.* 193 (1): 213–221
17. Schuchardt JP, Neubronner J, Kressel G, et al. (2011) Moderate doses of EPA and DHA from re-esterified triacylglycerols but not from ethyl-esters lower fasting serum triacylglycerols in statin-treated dyslipidemic subjects: Results from a six month randomized controlled trial. *Prostaglandins Leukot Essent Fatty Acids.* 85 (6): 381–386
18. Dyerberg J, Madsen P, Moller JM, et al. (2010) Bioavailability of marine n-3 fatty acid formulations. *Prostaglandins Leukot Essent Fatty Acids.* 83 (3): 137–141
19. Harris WS, De Caterina R, Marik PE (2013) Docosahexaenoic acid ethyl esters ineffective? *Proc Natl Acad Sci U S A.* 110 (25):E2259
20. Wu SY, Mayneris-Perxachs J, Lovegrove JA, et al. (2014) Fish-oil supplementation alters numbers of circulating endothelial progenitor cells and microparticles independently of eNOS genotype. *AM J CLIN NUTR.* 100 (5): 1232–1243
21. Cazzola R, Russo-Volpe S, Miles EA, et al. (2007) Age- and dose-dependent effects of an eicosapentaenoic acid-rich oil on cardiovascular risk factors in healthy male subjects. *ATHEROSCLEROSIS.* 193 (1): 159–167
22. Deutsche Gesellschaft für Ernährung e. V. (DGE) (2015) *Evidenzbasierte Leitlinie: „Fettzufuhr und Prävention ausgewählter ernährungsmitbedingter Krankheiten“, 2. Version* (2015).
23. Nelson RH (2013) Hyperlipidemia as a risk factor for cardiovascular disease. *PRIMARY CARE.* 40 (1):195–211
24. Browning LM, Krebs JD, Moore CS, et al. (2007) The impact of long chain n-3 polyunsaturated fatty acid supplementation on inflammation, insulin sensitivity and CVD risk in a group of overweight women with an inflammatory phenotype. *DIABETES OBES METAB.* 9 (1): 70–80
25. Goodfellow J, Bellamy MF, Ramsey MW, et al. (2000) Dietary supplementation with marine omega-3 fatty acids improve systemic large artery endothelial function in subjects with hypercholesterolemia. *J AM COLL CARDIOL.* 35 (2): 265–270
26. Hartweg J, Perera R, Montori V, et al. (2008) Omega-3 polyunsaturated fatty acids (PUFA) for type 2 diabetes mellitus. *COCHRANE DB SYST REV.* 1: CD003205
27. Ciubotaru, I, Lee YS, Wander RC (2003) Dietary fish oil decreases C-reactive protein, interleukin-6, and triacylglycerol to HDL-cholesterol ratio in postmenopausal women on HRT. *J NUTR BIOCHEM.* 14 (9): 513–521
28. Skulas-Ray AC, Kris-Etherton PM, Harris WS, et al. (2011) Dose-response effects of omega-3 fatty acids on triglycerides, inflammation, and endothelial function in healthy persons with moderate hypertriglyceridemia. *Am J Clin Nutr.* 93(2):243–252
29. Harris WS (1999) Nonpharmacologic Treatment of Hypertriglyceridemia: Focus on Fish Oils. *Clin. Cardiol.* 22(6 Suppl):II40-II43.
30. Jump DB, Botolin D, Wan Y, et al. (2008) Docosahexaenoic acid (DHA) and hepatic gene transcription. *CHEM PHYS LIPIDS.* 153 (1): 3–13
31. Grygiel-Gróniak B (2014) Peroxisome proliferator-activated receptors and their ligands: nutritional and clinical implications – a review. *Nutr J.* 13:17
32. Miller M, Langenberg P, Havas S (2007) Impact of lowering triglycerides on raising HDL-C in hypertriglyceridemic and non-hypertriglyceridemic subjects. *Int J Cardiol.* 119(2): 192–195
33. Jacobson GA, Glickstein SB, Rowe JD, et al. (2012) Effects of eicosapentaenoic acid and docosahexaenoic acid on low-density lipoprotein cholesterol and other lipids: a review. *J Clin Lipidol.* 6(1):5–18
34. Lankinen M, Kolehmainen M, Jaaskelainen T, et al. (2014) Effects of Whole Grain, Fish and Bilberries on Serum Metabolic Profile

- and Lipid Transfer Protein Activities: A Randomized Trial (Sysdimet). *PLOS ONE*. 9 (2)
35. Maki KC, Van Elswyk ME, Alexander DD, et al. (2012) A meta-analysis of randomized controlled trials that compare the lipid effects of beef versus poultry and/or fish consumption. *J Clin Lipidol*. 6 (4): 352–361
 36. Adamsson V, Cederholm T, Vessby B, et al. (2014) Influence of a healthy Nordic diet on serum fatty acid composition and associations with blood lipoproteins – results from the NORDIET study. *J FOOD NUTR RES*. 58: 24114
 37. Bulliyya G (2002) Influence of fish consumption on the distribution of serum cholesterol in lipoprotein fractions: comparative study among fish-consuming and non-fish-consuming populations. *ASIA PAC J CLIN NUTR*. 11(2): 104–111
 38. Calder PC (2006) n-3 polyunsaturated fatty acids, inflammation, and inflammatory diseases. *AM J CLIN NUTR*. 83 (6): 1505S–1519S
 39. Caterina R, Madonna R, Bertolotto A (2007) n-3 fatty acids in the treatment of diabetic patients – Biological rationale and clinical data. *DIABETES CARE*. 30 (4): 1012–1026
 40. Li K, Huang T, Zheng J (2014) Effect of marine-derived n-3 polyunsaturated fatty acids on C-reactive protein, interleukin 6 and tumor necrosis factor α : a meta-analysis. *PLOS ONE*. 9 (2): e88103
 41. Lopez-Garcia E, Schulze MB, Manson JE (2004) Consumption of (n-3) fatty acids is related to plasma biomarkers of inflammation and endothelial activation in women. *J NUTR*. 134 (7): 1806–1811
 42. Eschen O, Christensen JH, Caterina R, et al. (2004) Soluble adhesion molecules in healthy subjects: a dose-response study using n-3 fatty acids. *NUTR METAB CARDIOVAS*. 14 (4): 180–185
 43. Hjerkin EM, Seljeflot I, Ellingsen I, et al. (2005) Influence of long-term intervention with dietary counseling, long-chain n-3 fatty acid supplements, or both on circulating markers of endothelial activation in men with long-standing hyperlipidemia. *AM J CLIN NUTR*. 81 (3): 583–589
 44. Miles EA, Thies F, Wallace FA (2001) Influence of age and dietary fish oil on plasma soluble adhesion molecule concentrations. *CLIN SCI*. 100 (1):91–100
 45. Rees D, Miles EA, Banerjee T et al. (2006) Dose-related effects of eicosapentaenoic acid on innate immune function in healthy humans: a comparison of young and older men. *AM J CLIN NUTR*. 83 (2): 331–342
 46. Bolton-Smith C, Woodward M, Tavendale R (1997) Evidence for age-related differences in the fatty acid composition of human adipose tissue, independent of diet. *EUR J CLIN NUTR*. 51 (9): 619–624
 47. Jackson KG, Armah CK, Doman I, et al. (2009) The impact of age on the post-prandial vascular response to a fish oil-enriched meal. *BRIT J NUTR*. 102 (10): 1414–1419
 48. Geelen A, Brouwer IA, Schouten EG, et al. (2004) Intake of n-3 fatty acids from fish does not lower serum concentrations of C-reactive protein in healthy subjects. *EUR J CLIN NUTR* 58 (10):1440–1442
 49. Madsen T, Christensen JH, Blom M, et al. (2003) The effect of dietary n-3 fatty acids on serum concentrations of C-reactive protein: a dose-response study. *BRIT J NUTR*. 89 (4): 517–522
 50. Vega-López S, Kaul N, Devaraj S, et al. (2004) Supplementation with omega3 polyunsaturated fatty acids and all-rac alpha-tocopherol alone and in combination failed to exert an anti-inflammatory effect in human volunteers. *Metabolism*. 53 (2): 236–240
 51. Katan MB, Deslypere JP, vanBirtelen APJM, et al. (1997) Kinetics of the incorporation of dietary fatty acids into serum cholesteryl esters, erythrocyte membranes, and adipose tissue: an 18-month controlled study. *J LIPID RES*. 38 (10): 2012–2022
 52. Patel PS, Bingham SA, Sharp SJ, et al. (2009) Association Between Type of Dietary Fish and Seafood Intake and the Risk of Incident Type 2 Diabetes The European Prospective Investigation of Cancer (EPIC)-Norfolk cohort study. *Diabetes Care*. 32 (10): 1857–1863
 53. Sotos PM, Guillen M, Sorli JV, et al. (2011) Meat and fish consumption in a high cardiovascular risk spanish mediterranean population. *Nutrition Hospitalaria*. 26 (5): 1033–1040
 54. Djousse L, Gaziano JM, Buring JE, et al. (2011) Dietary omega-3 fatty acids and fish consumption and risk of type 2 diabetes. *AM J CLIN NUTR*. 93 (1): 143–150
 55. Ouellet V, Marois J, Weisnagel SJ, et al. (2007) Dietary cod protein improves insulin sensitivity in insulin-resistant men and women. *Diabetes Care*. 30 (11): 2816–2821
 56. Huang T, Wahlqvist ML, Xu T, et al. (2010) Increased plasma n-3 polyunsaturated fatty acid is associated with improved insulin sensitivity in type 2 diabetes in China. *MOL NUTR FOOD RES*. 54: S112–S119
 57. Woodman RJ, Mori TA, Burke V, et al. (2002) Effects of purified eicosapentaenoic and docosahexaenoic acids on glycemic control, blood pressure, and serum lipids in type 2 diabetic patients with treated hypertension. *AM J CLIN NUTR*. 76 (5): 1007–1015
 58. Muley A, Muley P, Shah M (2014) ALA, fatty fish or marine n-3 fatty acids for preventing DM?: a systematic review and meta-analysis. *Curr Diabetes Rev*. 10 (3): 158–165
 59. Wallin A, Di Giuseppe D, Orsini N, et al. (2012) Fish Consumption, Dietary Long-Chain n-3 Fatty Acids, and Risk of Type 2 Diabetes Systematic review and meta-analysis of prospective studies. *Diabetes Care*. 35 (4): 918–929
 60. Zhang M, Picard-Deland E, Marette A (2013) Fish and marine omega-3 polyunsaturated Fatty Acid consumption and incidence of type 2 diabetes: a systematic review and meta-analysis. *Int J Endocrinol*. 501015
 61. Zheng JS, Huang T, Yang J, et al. (2012) Marine N-3 polyunsaturated fatty acids are inversely associated with risk of type 2 diabetes in Asians: a systematic review and meta-analysis. *PLOS ONE*. 7 (9): e44525
 62. Chen R, Corona E, Sikora M, et al. (2012) Type 2 diabetes risk alleles demonstrate extreme directional differentiation among human populations, compared to other diseases. *PLoS genetics*. 8 (4): e1002621
 63. Schulze MB, Hoffmann K, Manson JE, et al. (2005) Dietary pattern, inflammation, and incidence of type 2 diabetes in women. *AM J CLIN NUTR*. 82 (3): 675–84
 64. Wall R, Ross RP, Fitzgerald GF, et al. (2010) Fatty acids from fish: the anti-inflammatory potential of long-chain omega-3 fatty acids. *NUTR REV*. 68 (5): 280–289
 65. World Cancer Research Fund International, American Institute for Cancer Research: Food, nutrition, physical activity, and the prevention of cancer: A global perspective. 1. publ. Washington, DC: AICR (2007)
 66. Shen XY, Zhou JD, Dong JY (2012) Dietary intake of n-3 fatty acids and colorectal cancer risk: a meta-analysis of data from

- 489 000 individuals. *BR J NUTR.* 108 (9): 1550–1556
67. Yu XF, Zou J, Dong J (2014) Fish consumption and risk of gastrointestinal cancers: A meta-analysis of cohort studies. *WORLD J GASTROENTERO.* 20 (41): 15398–15412
68. Liu J, Ma DWL (2014) The Role of n-3 Polyunsaturated Fatty Acids in the Prevention and Treatment of Breast Cancer. *Nutrients.* 6 (11): 5184–5223
69. Patterson RE, Flatt SW, Newman VA (2011) Marine Fatty Acid Intake Is Associated with Breast Cancer Prognosis. *J NUTR.* 141 (2): 201–206
70. Zheng JS, Hu XJ, Zhao YM, et al. (2013) Intake of fish and marine n-3 polyunsaturated fatty acids and risk of breast cancer: meta-analysis of data from 21 independent prospective cohort studies. *BRIT MED J.* 346
71. Saadatian-Elahi M, Norat T, Goudable J, et al. (2004) Biomarkers of dietary fatty acid intake and the risk of breast cancer: a meta-analysis. *INT J CANCER.* 111 (4): 584–591
72. Schmidt JA, Gorst-Rasmussen A, Nystrom PW, et al. (2014) Baseline patterns of adipose tissue fatty acids and long-term risk of breast cancer: a case-cohort study in the Danish cohort Diet, Cancer and Health. *EUR J CLIN NUTR.* 68 (10): 1088–1094
73. Witt PM, Christensen JH, Schmidt EB, et al. (2009) Marine n-3 polyunsaturated fatty acids in adipose tissue and breast cancer risk: a case-cohort study from Denmark. *CANCER CAUSE CONTROL.* 20 (9): 1715–1721
74. Amin M, Jeyaganth S, Fahmy N, et al. (2008) Dietary habits and prostate cancer detection: a case-control study. *CUAJ-CAN UROL ASSOC.* 2 (5): 510–515
75. Leitzmann MF, Stampfer MJ, Michaud DS, et al. (2004) Dietary intake of n-3 and n-6 fatty acids and the risk of prostate cancer. *AM J CLIN NUTR.* 80 (1): 204–216
76. Terry P, Lichtenstein P, Feychting M, et al. (2001) Fatty fish consumption and risk of prostate cancer. *Lancet.* 357 (9270): 1764–1766
77. Lovegrove C, Ahmed K, Challacombe B, et al. (2015) Systematic review of prostate cancer risk and association with consumption of fish and fish-oils: analysis of 495,321 participants. *INT J CLIN PRACT.* 69 (1): 87–105
78. Pham TM, Fujino Y, Kubo T, et al. (2009) Fish intake and the risk of fatal prostate cancer: findings from a cohort study in Japan. *PUBLIC HEALTH NUTR.* 12 (5): 609–613
79. Chavarro JE, Stampfer MJ, Hall MN, et al. (2008) A 22-y prospective study of fish intake in relation to prostate cancer incidence and mortality. *AM J CLIN NUTR.* 88 (5): 1297–1303
80. Alexander D, Bassett J, Weed D, et al. (2015) Meta-Analysis of Long-Chain Omega-3 Polyunsaturated Fatty Acids (LCv-3PUFA) and Prostate Cancer. *NUTR CANCER.* 67(4), 543–554
81. Chua ME, Sio MC, Sorongon MC et al. (2012) Relationship of dietary intake of omega-3 and omega-6 fatty acids with risk of prostate cancer development: a meta-analysis of prospective studies and review of literature. *PROSTATE CANCER.* 2012: 826254.
82. Scientific Opinion on the Tolerable Upper Intake Level of eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) and docosapentaenoic acid (DPA) : European Food Safety Authority (EFSA). (2012)
83. Nationale Verzehrs-Studie II. [Karlsruhe] : Max Rubner Institut, Bundesforschungsinstitut für Ernährung und Lebensmittel
84. Lederman SA, Jones RL, Caldwell KL, et al. (2008) Relation between cord blood mercury levels and early child development in a World Trade Center cohort. *ENVIRON HEALTH PERSP.* 116 (8): 1085–1091
85. Mahaffey KR, Sunderland EM, Chan HM, et al. (2011): Balancing the benefits of n-3 polyunsaturated fatty acids and the risks of methylmercury exposure from fish consumption. *NUTR REV.* 69 (9): 493–508
86. Bundesinstitut für Risikobewertung: Aufnahme von Umweltkontaminanten über Lebensmittel Ergebnisse des Forschungsprojektes LEXUKon. URL http://www.bfr.bund.de/cm/350/aufnahme_von_umweltkontaminanten_ueber_lebensmittel.pdf – Zugriff: 11.01.15
87. European Food Safety Authority (EFSA) (2015): Statement on the benefits of fish/seafood consumption compared to the risks of methylmercury in fish/seafood. *EFSA JOURNAL.* 13(1):3982
88. Bundesinstitut für Risikobewertung: Quecksilber und Methylquecksilber in Fischen und Fischprodukten – Bewertung durch die EFSA : Stellungnahme des BfR vom 29. März 2004. URL http://www.bfr.bund.de/cm/343/quecksilber_und_methylquecksilber_in_fischen_und_fischprodukten__bewertung_durch_die_efs.pdf – Zugriff: 20.01.2015

DOI: 10.4455/eu.2015.026