Literatur

- Wolfram G, Bechthold A, Boeing H et al. (2015) Evidence-based guideline of the German Nutrition Society: fat intake and prevention of selected nutrition-related diseases. Ann Nutr Metab 67: 141–204
- 2. Hooper L, Martin N, Abdelhamid A, Davey Smith G (2015) Reduction in saturated fat intake for cardiovascular disease. Cochrane Database Syst Rev(6): Cd011737
- 3. Mozaffarian D, Micha R, Wallace S (2010) Effects on coronary heart disease of increasing polyunsaturated fat in place of saturated fat: a systematic review and meta-analysis of randomized controlled trials. PLoS Med 7: e1000252
- Ramsden CE, Zamora D, Majchrzak-Hong S et al. (2016) Re-evaluation of the traditional diet-heart hypothesis: analysis of recovered data from minnesota coronary experiment (1968–73). BMJ 353: i1246
- Ramsden CE, Zamora D, Leelarthaepin B et al. (2013) Use of dietary linoleic acid for secondary prevention of coronary heart disease and death: evaluation of recovered data from the Sydney diet heart study and updated metaanalysis. BMJ 346: e8707
- Chowdhury R, Warnakula S, Kunutsor S et al. (2014) Association of dietary, circulating, and supplement fatty acids with coronary risk: a systematic review and meta-analysis. Ann Intern Med 160: 398–406
- de Souza RJ, Mente A, Maroleanu A et al. (2015) Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies. BMJ 351: h3978
- 8. Li Y, Hruby A, Bernstein AM et al. (2015) Saturated fats compared with unsaturated fats and sources of carbohydrates in relation to risk of coronary heart disease: a prospective cohort study. J Am Coll Cardiol 66: 1538–1548
- Estruch R, Ros E, Salas-Salvado J et al. (2013) Primary prevention of cardiovascular disease with a mediterranean diet. N Engl J Med 368: 1279–1290
- 10. Lopez-Garcia E, Rodriguez-Artalejo F, Li TY et al. (2014) The mediterranean-style dietary pattern and mortality among men and women with cardiovascular disease. Am J Clin Nutr 99: 172–180
- 11. Fung TT, Rexrode KM, Mantzoros CS et al. (2009) Mediterranean diet and incidence of and mortality from coronary heart disease and stroke in women. Circulation 119: 1093–1100

- Mozaffarian D, Lemaitre RN, King IB et al. (2013) Plasma phospholipid long-chain omega-3 fatty acids and total and causespecific mortality in older adults: A cohort study. Ann Intern Med 158: 515–525
- 13. de Oliveira Otto MC, Wu JH, Baylin A et al. (2013) Circulating and dietary omega-3 and omega-6 polyunsaturated fatty acids and incidence of cvd in the multi-ethnic study of atherosclerosis. J Am Heart Assoc 2: e000506
- 14. Kleber ME, Delgado GE, Lorkowski S et al. (2016) Omega-3 fatty acids and mortality in patients referred for coronary angiography. The Ludwigshafen risk and cardiovascular health study. Atherosclerosis 252: 175–181
- Dawczynski C, Lorkowski S (2016) Transfatty acids and cardiovascular risk: does origin matter? Expert Rev Cardiovasc Ther 14: 1001–1005
- Bloomfield HE, Koeller E, Greer N et al. (2016) Effects on health outcomes of a mediterranean diet with no restriction on fat intake: a systematic review and meta-analysis. Ann Intern Med 165: 491–500
- 17. Smart NA, Marshall BJ, Daley M et al. (2011) Low-fat diets for acquired hypercholesterolaemia. Cochrane Database Syst Rev (2): Cd007957
- 18. St-Onge MP, Bosarge A, Goree LL, Darnell B (2008) Medium chain triglyceride oil consumption as part of a weight loss diet does not lead to an adverse metabolic profile when compared to olive oil. J Am Coll Nutr 27: 547–552
- Nagao K, Yanagita T (2010) Medium-chain fatty acids: functional lipids for the prevention and treatment of the metabolic syndrome. Pharmacol Res 61: 208–212
- U.S. Department of Health and Human Services, U.S. Department of Agriculture. Dietary guidelines for Americans 2015–2010, ed 8. (2015)
- 21. Mozaffarian D, Ludwig DS (2015) The 2015 us dietary guidelines: lifting the ban on total dietary fat. JAMA 313: 2421–2422
- 22. Mozaffarian D, Ludwig DS (2015) Dietary cholesterol and blood cholesterol contrations (reply). JAMA 314: 2084–2085
- 23. Rong Y, Chen L, Zhu T et al. (2013) Egg consumption and risk of coronary heart disease and stroke: Dose-response meta-analysis of prospective cohort studies. BMJ 346: e8539
- 24. Levin S, Wells C, Barnard N (2015) Dietary cholesterol and blood cholesterol concentrations. JAMA 314: 2083–2084

- Berger S, Raman G, Vishwanathan R et al. (2015) Dietary cholesterol and cardiovascular disease: a systematic review and meta-analysis. Am J Clin Nutr 102: 276–294
- 26. Warensjo E, Jansson JH, Berglund L et al. (2004) Estimated intake of milk fat is negatively associated with cardiovascular risk factors and does not increase the risk of a first acute myocardial infarction. A prospective case-control study. Br J Nutr 91: 635–642
- 27. Warensjo E, Jansson JH, Cederholm T et al. (2010) Biomarkers of milk fat and the risk of myocardial infarction in men and women: A prospective, matched case-control study. Am J Clin Nutr 92: 194–202
- 28. de Oliveira Otto MC, Mozaffarian D, Kromhout D et al. (2012) Dietary intake of saturated fat by food source and incident cardiovascular disease: The multi-ethnic study of atherosclerosis. Am J Clin Nutr 96: 397–404
- 29. Hussey EK, Portelli S, Fossler MJ et al. (2012) Relative bioavailability of an emulsion formulation for omega-3-acid ethyl esters compared to the commercially available formulation: A randomized, parallel-group, single-dose study followed by repeat dosing in healthy volunteers. Clin Pharmacol Drug Dev 1: 14–23
- 30. Davidson MH, Johnson J, Rooney MW et al. (2012) A novel omega-3 free fatty acid formulation has dramatically improved bioavailability during a low-fat diet compared with omega-3-acid ethyl esters: The eclipse (epanova® compared to lovaza®) in a pharmacokinetic single-dose evaluation) study. J Clin Lipidol 6: 573–584
- Schonfeld G, Patsch W, Rudel LL et al. (1982) Effects of dietary cholesterol and fatty acids on plasma lipoproteins. J Clin Invest 69: 1072– 1080