

Literatur

1. DGKJ (2013) Vitamin-K-Prophylaxe bei Neugeborenen. Empfehlungen der Ernährungskommission der Deutschen Gesellschaft für Kinder- und Jugendmedizin (DGKJ). *Monatsschr Kinderheilkd* 161: 351–353
2. DGE, ÖGE, SGE. D-A-CH-Referenzwerte für die Nährstoffzufuhr. 2. Aufl., Neuer Umschau Buchverlag, Neustadt a. d. Weinstraße (2015)
3. Nobel Media (2014) Edward A Doisy – Biographical. URL: www.nobelprize.org/nobel_prizes/medicine/laureates/1943/doisy-bio.html Zugriff 29.01.17
4. Nobel Media (2014) Henrik Dam – Biographical. URL: www.nobelprize.org/nobel_prizes/medicine/laureates/1943/dam-bio.html Zugriff 29.01.17
5. Schurgers LJ, Vermeer C (2000) Determination of phylloquinone and menaquinones in food. *Haemostasis* 30: 298–307
6. Booth SL (2002) Vitamin K: food composition and dietary intakes. *Food Nutr Res* 56: 5505
7. Beulens JW, Booth SL, van der Heuvel EG et al. (2013) The role of menaquinones (vitamin K2) in human health. *Br J Nutr* 110: 1357–1368
8. Nimptsch K, Rohrmann S, Linseisen J (2008) Dietary intake of vitamin K and risk of prostate cancer in the Heidelberg cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC Heidelberg). *Am J Clin Nutr* 87: 985–992
9. Beulens JW, van der A DL, Grobbee DE et al. (2010) Dietary phylloquinone and menaquinones intakes and risk of type 2 diabetes. *Diabetes Care* 33: 1699–1705
10. Schurgers LJ, Teunissen KJ, Hamulyá K et al. (2007) Vitamin K-containing dietary supplements: comparison of synthetic vitamin K1 and natto-derived menaquinone-7. *Blood* 109: 3279–3283
11. Olson RE (1984) The function and metabolism of vitamin K. *Annu Rev Nutr* 4: 281–337
12. EFSA (2017) Dietary reference values for vitamin K. *EFSA Journal* [in Vorbereitung]. URL: www.efsa.europa.eu/sites/default/files/consultation/170113.pdf Zugriff 20.01.17
13. Thijssen HHW, Vervoort LMT, Schurgers LJ et al. (2006) Menadiol is a metabolite of oral vitamin K. *Br J Nutr* 95: 260–266
14. Westenfeld R, Krueger T, Schlieper G et al. (2012) Effect of vitamin K2 supplementation on functional vitamin K deficiency in hemodialysis patients: a randomized trial. *Am J Kidney Dis* 59: 186–195
15. Schurgers LJ, Vermeer C (2002) Differential lipoprotein transport pathways of K-vitamins in healthy subjects. *Biochim Biophys Acta* 1570: 27–32
16. Vermeer C (2012) Vitamin K: the effect on health beyond coagulation – an overview. *Food Nutr Res* 56: 5329
17. Sato T (2012) Vitamin K2 and bone quality. *Vitam Trace Elem S6*: 001
18. Shea MK, Booth SL (2016) Concepts and controversies in evaluating vitamin K status in population-based studies. *Nutrients* 8: E8
19. Booth SL, Martini L, Peterson JW et al. (2003) Dietary phylloquinone depletion and repletion in older women. *J Nutr* 133: 2565–2569
20. Sato T, Schurgers LJ, Uenishi K (2012) Comparison of menaquinone-4 and menaquinone-7 on bioavailability in healthy women. *Nutr J* 11: 93
21. Olson RE, Chao J, Graham D et al. (2007) Total body phylloquinone and its turnover in human subjects at two levels of vitamin K intake. *Br J Nutr* 87: 543–553
22. EFSA (2006) Tolerable upper intake levels of vitamins and minerals. URL: www.efsa.europa.eu/sites/default/files/efsa_rep/blob_server_assets/ndatolerableuil.pdf Zugriff 31.01.17
23. Institute of Medicine. Dietary reference intakes. The essential guide to nutrient requirements. National Academic Press, Washington (2006)
24. McKeown NM, Jacques PF, Gundberg CM et al. (2002) Dietary and nondietary determinants of vitamin K biochemical measures in men and women. *J Nutr* 132: 1329–1334
25. Geleijnse JM, Vermeer C, Grobbee DE et al. (2004) Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: The Rotterdam Study. *J Nutr* 134: 3100–3105
26. Vissers LET, Dalmeijer GW, Boer JMA et al. (2013) Intake of dietary phylloquinone and menaquinones and risk of stroke. *J Am Heart Assoc* 2(6): e000455
27. Gast GC, de Roos NM, Sluijs I et al. (2009) A high menaquinone intake reduces the incidence of coronary heart disease. *Nutr Metab Cardiovasc Dis* 19: 504–510
28. Mensink GBM, Heseke H, Richter A et al. (2007) Ernährungsstudie als KIGGS-Modul (EsKiMo). URL: www.bmel.de/SharedDocs/Downloads/Ernaehrung/EsKiMoStudie.pdf?__blob=publicationFile Zugriff 31.01.17
29. Office for National Statistics. Social and Vital Statistics Division, Food Standards Agency (2005) National Diet and Nutrition Survey: adults aged 19 to 64 years, 2000–2001. [data collection]. UK Data Service. SN: 5140
30. EFSA (2008) Vitamin K2 added for nutritional purposes in foods for particular nutritional uses, food supplements and foods intended for the general population and Vitamin K2 as a source of vitamin K added for nutritional purposes to foodstuffs, in the context of regulation (EC) N° 258/97. *EFSA Journal* 6: 1–31
31. RIVM (2016) MEMO: Inname van vitamine K door kinderen en volwassenen in Nederland. Resultaten van VCP 2007–2010. URL: www.rivm.nl/dsresource?objectid=b96a6448-882a-41c1-bb72-6ece306bc4b2&type=org&disposition=inline Zugriff 31.01.17
32. DGE. 12. Ernährungsbericht. Bonn (2012)
33. SCF (1993) Nutrient and energy intakes for the European community food – science and technique. Reports of the Scientific Committee for Food (31st Series). Brüssel: Commission of the European Communities. URL: https://ec.europa.eu/food/sites/food/files/safety/docs/sci-com_scf_out89.pdf Zugriff 20.01.17
34. Booth SL, Dallal G, Shea MK et al. (2008) Effect of vitamin K supplementation on bone loss in elderly men and women. *J Clin Endocrinol Metab* 93: 1217–1223
35. Cranenburg EC, Koos R, Schurgers LJ et al. (2010) Characterisation and potential diagnostic value of circulating matrix gla protein (MGP) species. *Thromb Haemost* 104: 811–822
36. Gundberg CM, Lian JB, Booth SL (2012) Vitamin K-dependent carboxylation of osteocalcin: friend or foe? *Adv Nutr* 3: 149–157
37. McCann JC, Ames BN (2009) Vitamin K, an example of triage theory: is micronutrient inadequacy linked to diseases of aging? *Am J Clin Nutr* 90: 889–907
38. Theuwissen E, Cranenburg EC, Knapen MH et al. (2012) Low-dose menaquinone-7 supplementation improved extra-hepatic vitamin K status, but had no effect on thrombin

- generation in healthy subjects. *Br J Nutr* 108: 1652–1657
39. Theuwissen E, Magdeleyns EJ, Braam LAJ et al. (2014) Vitamin K status in healthy volunteers. *Food Funct* 5: 229–234
40. Dowd P, Ham SW, Geib SJ (1991) Mechanism of action of vitamin K. *J Am Chem Soc* 113: 7734–7743
41. Tie J-K, Jin D-Y, Straight DL et al. (2011) Functional study of the vitamin K cycle in mammalian cells. *Blood* 117: 2967–2974
42. van den Heuvel EG, van Schoor NM, Lips P et al. (2014) Circulating uncarboxylated matrix Gla protein, a marker of vitamin K status, as a risk factor for cardiovascular diseases. *Maturitas* 77: 137–141
43. Theuwissen E, Teunissen KJ, Spronk HM et al. (2013) Effect of low-dose supplements of menaquinone-7 (vitamin K₂) on the stability of oral anticoagulant treatment: dose-response relationship in healthy volunteers. *J Thromb Haemost* 11: 1085–1092
44. Viegas CSB, Simes DC (2016) New perspectives for the nutritional value of vitamin K in human health. *J Nutr Disorders Ther* 6: 192
45. Barnes C, Newall F, Ignjatovic V et al. (2005) Reduced bone density in children on long-term warfarin. *Pediatr Res* 57: 578–581
46. Rezaieyazdi Z, Falsoleiman H, Khajehdaluee M et al. (2009) Reduced bone density in patients on long-term warfarin. *Int J Rheum Dis* 12: 130–135
47. ML, Winkers K, Hackeng TM et al. (2012) Vascular calcification: the prize to pay for anticoagulation therapy with vitamin K-antagonists. *Blood Rev* 26: 155–166
48. Rennenberg RJMW, van Varik BJ, Schurgers LJ et al. (2010) Chronic coumarin treatment is associated with increased extracoronary arterial calcifications in humans. *Blood* 115: 5121–5123
49. Schurgers LJ, Joosen IA, Laufer EM et al. (2012) Vitamin-K-antagonists accelerate atherosclerotic calcification and induce a vulnerable plaque phenotype. *PloS One* 7: e43229
50. Karlson B, Leijd B, Hellström K (1986) On the influence of vitamin K-rich vegetables and wine on the effectiveness of warfarin treatment. *Acta Med Scand* 220: 347–350
51. Violi F, Lip GYH, Pignatelli P et al. (2016) Interaction between dietary vitamin K intake and anticoagulation by vitamin K antagonists: is it really true? A systematic review. *Medicine (Baltimore)* 95: e2895
52. Rasmussen MA, Skov J, Bladbjerg E-M et al. (2012) Multivariate analysis of the relation between diet and warfarin dose. *Eur J Clin Pharmacol* 68: 321–328
53. Schurgers LJ, Shearer MJ, Hamulyák K et al. (2004) Effect of vitamin K intake on the stability of oral anticoagulant treatment: dose-response relationships in healthy subjects. *Blood* 104: 2682–2689
54. Chatrou MLL, Reutelingsperger CR, Schurgers LJ (2011) Role of vitamin K dependent proteins in the arterial vessel wall. *Hämostaseologie* 31: 251–257
55. Theuwissen E, Smit E, Vermeer C (2012) The role of vitamin K in soft-tissue calcification. *Adv Nutr* 3: 166–173
56. Viegas CSB, Simes DC (2016) Gla-rich protein (GRP): a new player in the burden of vascular calcification. *J Cardiovasc Dis Diagn* 4: 245
57. Nakajima S, Iijima H, Egawa S et al. (2011) Association of vitamin K deficiency with bone metabolism and clinical disease activity in inflammatory bowel disease. *Nutrition* 27: 1023–1028
58. Schoon EJ, Müller MC, Vermeer C et al. (2001) Low serum and bone vitamin K status in patients with longstanding Crohn's disease: another pathogenetic factor of osteoporosis in Crohn's disease? *Gut* 48: 473–477
59. Booth SL, Golly I, Sackeck JM et al. (2004) Effect of vitamin E supplementation on vitamin K status in adults with normal coagulation status. *Am J Clin Nutr* 80: 143–148
60. Fusaro M, Noale M, Viola V et al. (2012) Vitamin K, vertebral fractures, vascular calcifications, an mortality: Vitamin K Italian (VIKI) Dialysis study. *J Bone Min Res* 27: 2271–2278
61. Persy V, D'Haese P (2009) Vascular calcification and bone disease: the calcification paradox. *Trends Mol Med* 15: 405–416
62. Rhéaume-Bleue K. Vitamin K₂ and the calcium paradox: how a little-known vitamin could save your life. Wiley & Sons Canada, Mississauga (2013)
63. Schek A (2017) Aktuelles in Kürze: Vitamin D – wie viel ist genug? *Leistungssport* 47: 25–27
64. Schek A. *Ernährungslehre kompakt*. 6. Aufl., Umschau Zeitschriftenverlag, Wiesbaden (2017)
65. Bolland MJ, Avenell A, Baron JA et al. (2010) Effect of calcium supplements on risk of myocardial infarction and cardiovascular events: meta-analysis. *BMJ* 341: c3691
66. Kanellakis S, Moschonis G, Tenta R et al. (2012) Changes in parameters of bone metabolism in postmenopausal women following a 12-month intervention period using dairy products enriched with calcium, vitamin D, and phylloquinone (vitamin K₁) or menaquinone-7 (vitamin K₂): the Postmenopausal Health Study II. *Calcif Tissue Int* 90: 251–262
67. Booth SL, Al Ragabi A (2008) Determinants of vitamin K status in humans. *Vitam Horm* 78: 1–22
68. Kaneki M, Hodges SJ, Hosoi T et al. (2001) Japanese fermented soybean food as the major determinant of the large geographic difference in circulating levels of vitamin K₂: possible implications for hip-fracture risk. *Nutrition* 17: 315–321
69. Tsugawa N, Shiraki M, Suhara Y et al. (2006) Vitamin K status of healthy Japanese women: age related vitamin K requirement for γ -carboxylation of osteocalcin. *Am J Clin Nutr* 83: 380–386
70. Tsugawa N, Shiraki M, Suhara Y et al. (2008) Low plasma phylloquinone concentration is associated with high incidence of vertebral fracture in Japanese women. *J Bone Miner Metab* 26: 79–85
71. Booth SL, Centi A, Smith SR et al. (2013) The role of osteocalcin in human glucose metabolism: marker oder mediator. *Nat Rev Endokrinol* 9: 43–55
72. Schurgers LJ, Uitto J, Reutelingsperger CP (2013) Vitamin K dependent carboxylation of matrix Gla-protein: a crucial switch to control ectopic mineralization. *Trends Mol Med* 19: 217–226
73. Mayer O, Seidlerová J, Bruthans J et al. (2014) Dephospho-uncarboxylated matrix Gla-protein is associated with mortality risk in patients with chronic stable vascular disease. *Atherosclerosis* 235: 162–168
74. van Summeren MJ, Vermeer C, Engelbert RH et al. (2008) Extremes in vitamin K status of bone are related to bone ultrasound properties in children with juvenile idiopathic arthritis. *Clin Exp Rheumatol* 26: 484–491
75. Villafán-Bernal JR, Llamas-Covarrubias MA, Muñoz-Valle JF et al. (2014) A cut-point value of uncarboxylated to carboxylated index is associated with glycemic status markers in type 2 diabetes. *J Investig Med* 62: 33–36
76. Schurgers LJ, Barreto DV, Barreto FC et al. (2010) The circulating inactive form of matrix gla protein is a surrogate marker of vascular calcification in chronic kidney disease:

- a preliminary report. *Clin J Am Soc Nephrol* 5: 568–575
77. Ueland T, Gullestad L, Dahl CP et al. (2010) Undercarboxylated matrix Gla protein is associated with indices of heart failure and mortality in symptomatic aortic stenosis. *J Intern Med* 268: 483–492
 78. Ueland T, Dahl CP, Gullestad I et al. (2011) Circulating levels of non-phosphorylated undercarboxylated matrix Gla protein are associated with disease severity in patients with chronic heart failure. *Clin Sci (Lond)* 121: 119–127
 79. Dalmeijer GW, van der Schouw YT, Vermeer C et al. (2013) Circulating matrix Gla protein is associated with coronary artery calcification and vitamin K status in healthy women. *J Nutr Biochem* 24: 624–628
 80. van Summeren MJ, Braam LA, Lilien MR et al. (2009) The effect of menaquinone-7 (vitamin K2) supplementation on osteocalcin carboxylation in healthy prepuberal children. *Br J Nutr* 102: 1171–1178
 81. Brugè F, Bacchetti C, Principi F et al. (2011) Olive oil supplemented with menaquinone-7 significantly increases osteocalcin carboxylation. *Br J Nutr* 106: 1058–1062
 82. Dalmeijer GW, van der Schouw YT, Magdeleyns E et al. (2012) The effect of menaquinone-7 supplementation on circulating species of matrix Gla protein. *Atherosclerosis* 225: 397–402
 83. Caluwé R, Vandecasteele S, van Vlem B et al. (2014) Vitamin K2 supplementation in haemodialysis patients: a randomized dose-finding study. *Nephrol Dial Transplant* 29: 1385–1390
 84. Knapen MH, Braam LA, Teunissen KJ et al. (2015) Yogurt drink fortified with menaquinone-7 improves vitamin K status in healthy population. *J Nutr Sci* 4: e35
 85. Kurnatowska I, Grzelak P, Masajtis-Zagajewka A et al. (2015) Effect of vitamin K2 on progression of atherosclerosis and vascular calcification in nondialyzed patients with chronic kidney disease stages 3–5. *Pol Arch Med Wevn* 125: 631–640
 86. van Gorp RH, Schurgers LJ (2015) New insights into the pros and cons of the clinical use of vitamin K antagonists (VKAs) versus direct oral anticoagulants (DOACs). *Nutrients* 7: 9538–9557
 87. Erkkilä AT, Booth SL, Hu FB et al. (2007) Phylloquinone intake and risk of cardiovascular diseases in men. *Nutr Metab Cardiovasc Dis* 17: 58–62
 88. Erkkilä AT, Booth SL, Hu FB et al. (2005) Phylloquinone intake as a marker for coronary heart disease risk but not stroke in women. *Eur J Clin Nutr* 59: 196–204
 89. Villines TC, Hatzigeorgiou C, Feuerstein IM et al. (2005) Vitamin K₁ intake and coronary calcification. *Coron Artery Dis* 16: 199–203
 90. Shea MK, O'Donnell CJ, Hoffmann U et al. (2009) Vitamin K supplementation and progression of coronary artery calcium in older men and women. *Am J Clin Nutr* 89: 1799–1807
 91. Beulens JW, Bots ML, Atsma F et al. (2009) High dietary menaquinone intake is associated with reduced coronary calcification. *Atherosclerosis* 203: 489–493
 92. Vissers LET, Dalmeijer GW, Boer JMA et al. (2016) The relationship between vitamin K and peripheral artery disease. *Atherosclerosis* 252: 15–20
 93. Knapen MH, Braam LA, Drummen ME et al. (2015) Menaquinone-7 supplementation improves arterial stiffness in healthy postmenopausal women: double-blind randomised clinical trial. *Thromb Haemost* 113: 1135–1144
 94. Vossen LM, Schurgers LJ, van Varik BJ et al. (2015) Menaquinone-7 supplementation to reduce vascular calcification in patients with coronary artery disease: rationale and study protocol (VitaK-CAC-Trial). *Nutrients* 7: 8905–8915
 95. Cauly JA (2015) Estrogen in bone health in men and women. *Steroids* 99(Pt A): 11–15
 96. Komarova SV, Safrank L, Gopalakrishnan J et al. (2015) Mathematical model for bone mineralization. *Front Cell Dev Biol* 3: 51
 97. Chen L, Jacquet R, Lowder E et al. (2015) Refinement of collagen-mineral interaction: a possible role for osteocalcin in apatite crystal nucleation, growth and development. *Bone* 71: 7–16
 98. Florencio-Silva R, Rodrigues da Silva Sasso G, Sasso-Cerri E et al. (2015) Biology of bone tissue: structure, function, and factors that influence bone cells. *Biomed Res Int* 2015: 421746
 99. Knapen MH, Schurgers LJ, Vermeer C (2007) Vitamin K₂ supplementation improves hip bone geometry and bone strength indices in postmenopausal women. *Osteoporos Int* 18: 963–972
 100. Plaza SM, Lamson DW (2005) Vitamin K₂ in bone metabolism and osteoporosis. *Altern Med Rev* 10: 24–35
 101. Yamaguchi M, Weitzmann MN (2011) Vitamin K2 stimulates osteoblastogenesis and suppresses osteoclastogenesis by suppressing NF-κB activation. *Int J Mol Med* 27: 3–14
 102. Cheung AM, Tile L, Lee Y et al. (2008) Vitamin K supplementation in postmenopausal women with osteopenia (ECKO trial): a randomized controlled trial. *PLoS Med* 5: 1–12
 103. Feskanich D, Weber P, Willett WC et al. (1999) Vitamin K intake and hip fractures in women: a prospective study. *Am J Clin Nutr* 69: 74–79
 104. Chan R, Leung J, Woo J (2012) No association between dietary vitamin K intake and fracture risk in Chinese community-dwelling older men and women: a prospective study. *Calcif Tissue Int* 90: 396–403
 105. Apalset EM, Gjesdal CG, Eide GE et al. (2011) Intake of vitamin K1 and K2 and risk of hip fractures: The Hordaland Health Study. *Bone* 49: 900–905
 106. Booth SL, Tucker KL, Chen H et al. (2000) Dietary vitamin K intakes are associated with hip fracture but not with bone mineral density in elderly men and women. *Am J Clin Nutr* 71: 1201–1208
 107. Booth SL, Broe KE, Gagnon DR et al. (2003) Vitamin K intake and bone mineral density in women and men. *Am J Clin Nutr* 77: 512–516
 108. Rejnmark L, Vestergaard P, Charles P et al. (2006) No effect of vitamin K₁ intake on bone mineral density and fracture risk in perimenopausal women. *Osteoporos Int* 17: 1122–1132
 109. Macdonald HM, McGuigan FE, Latham-New SA et al. (2008) Vitamin K1 intake is associated with higher bone mineral density and reduced bone resorption in early postmenopausal Scottish women: no evidence of gene-nutrient interaction with apolipoprotein E polymorphisms. *Am J Clin Nutr* 8: 1513–1520
 110. Bulló M, Estruch R, Salas-Salvadó J (2011) Dietary vitamin K intake is associated with bone quantitative ultrasound measurements but not with bone peripheral biochemical markers in elderly men and women. *Bone* 48: 1313–1318
 111. Volpe SL, Leung MM, Giordano H (2008) Vitamin K supplementation does not significantly impact bone mineral density and biochemical markers of bone in pre- and perimenopausal women. *Nutr Res* 28: 577–582
 112. Fujita Y, Iki M, Tamaki J et al. (2012) Association between vitamin K intake from fermented soybeans, natto, and bone mineral density in elderly Japanese men: the Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) study. *Osteoporos Int* 23: 705–714

113. Ikeda Y, Iki M, Morita A et al. (2006) Intake of fermented soybeans, natto, is associated with reduced bone loss in postmenopausal women: Japanese population-based osteoporosis (JPOS) study. *J Nutr* 136: 1323–1328
114. Knapen MH, Drummen NE, Smit E et al. (2013) Three-year low-dose menaquinone-7 supplementation helps decrease bone loss in healthy postmenopausal women. *Osteoporos Int* 24: 2499–2507
115. Emaus N, Gjesdal CG, Almas B et al. (2010) Vitamin K2 supplementation does not influence bone loss in early menopausal women: a randomised double-blind placebo-controlled trial. *Osteoporos Int* 21: 1731–1740
116. Binkley N, Harke J, Krueger D et al. (2009) Vitamin K treatment reduces undercarboxylated osteocalcin but does not alter bone turnover, density, or geometry in healthy postmenopausal North American women. *J Bone Min Res* 24: 983–991
117. Orimo H, Shiraki M, Tomita A et al. (1998) Effects of menatetrenone on the bone and calcium metabolism in osteoporosis: a double-blind placebo-controlled study. *J Bone Mineral Metab* 16: 116–112
118. Cockayne S, Adamson J, Lanham-New S et al. (2006) Vitamin K and the prevention of fractures. Systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med* 166: 1256–1261
119. Huang Z-B, Wan S-L, Lu Y-J et al. (2015) Does vitamin K2 play a role in the prevention and treatment of osteoporosis for postmenopausal women: a meta-analysis of randomized controlled trials. *Osteoporos Int* 26: 1175–1186
120. Fang Y, Hu C, Tao X et al. (2012) Effect of vitamin K on bone mineral density: a meta-analysis of randomized controlled trials. *J Bone Mineral Metab* 30: 60–68
121. Hamidi MS, Gajic-Veljanoski O, Cheung AM (2013) Vitamin K and bone health. *J Clin Densitom* 16: 409–413
122. Cavaco S, Viegas CSB, Rafael MS et al. (2015) Gla-rich protein is involved in the cross-talk between calcification and inflammation in osteoarthritis. *Cell Mol Life Sci* 73: 1051–1065
123. Oka H, Akune T, Muraki S et al. (2009) Association of low dietary vitamin K intake with radiographic knee osteoarthritis in Japanese elderly population: dietary survey in a population-based cohort of the ROAD study. *J Orthop Sci* 14: 687–692
124. Neogi T, Booth SL, Zhang YQ et al. (2006) Low vitamin K status is associated with osteoarthritis in the hand and knee. *Arthritis Rheum* 54: 1255–1262
125. Misra D, Booth SL, Tolstykh I et al. (2013) Vitamin K deficiency is associated with incident knee osteoarthritis. *Am J Med* 126: 243–248
126. Ferron M, Hinoi E, Karsenty G et al. (2008) Osteocalcin differentially regulates β cell and adipocyte gene expression and affects the development of metabolic diseases in wild-type mice. *Proc Natl Acad Sci USA* 105: 5266–5270
127. Choi HJ, Yu J, An JH et al. (2011) Vitamin K2 supplementation improves insulin sensitivity via osteocalcin metabolism: a placebo-controlled trial. *Diabetes Care* 34: e147
128. Yoshida M, Booth SL, Meigs JB et al. (2008) Phylloquinone intake, insulin sensitivity, and glycemic status in men and women. *Am J Clin Nutr* 88: 210–215
129. Kumar R, Binkley N, Vella A (2010) Effect of phylloquinone supplementation on glucose homeostasis in humans. *Am J Clin Nutr* 92: 1528–1532
130. Yoshida M, Jacques PF, Meigs JB et al. (2008) Effect of vitamin K supplementation on insulin resistance in older men and women. *Diabetes Care* 31: 2092–2096
131. Juanola-Falgarona M, Salas-Salvadó J, Estruch R et al. (2013) Association between dietary phylloquinone intake and peripheral metabolic risk markers related to insulin resistance and diabetes in elderly subjects at high cardiovascular risk. *Cardiovasc Diabetol* 12: 7
132. Zhang W, Shen X, Wan C et al. (2012) Effect of insulin and insulin-like growth factor 1 on osteoblast proliferation and differentiation: differential signalling via Akt and ERK. *Cell Biochem Funct* 30: 297–302
133. Janghorbani M, van Dam RM, Willett WC et al. (2007) Systematic review of type 1 and type 2 diabetes mellitus and risk of fracture. *Am J Epidemiol* 166: 495–505
134. Dam V, Dalmeijer GW, Vermeer C et al. (2015) Association between vitamin K and metabolic syndrome: a 10-year follow-up study in adults. *J Clin Endocrinol Metab* 100: 2472–2479
135. Viegas CS, Herfs M, Rafael MS et al. (2014) Gla-rich protein is a potential new vitamin K target in cancer. Evidences for a direct GPR-mineral interaction. *Biomed Res Int* 2014: 340216
136. Juanola-Falgarona M, Salas-Salvadó J, Martínez-González MA et al. (2014) Dietary intake of vitamin K is inversely associated with mortality risk. *J Nutr* 144: 743–750
137. Nimptsch K, Rohrmann S, Kaaks R et al. (2010) Dietary vitamin K intake in relation to cancer incidence and mortality: results from the Heidelberg cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Heidelberg). *Am J Clin Nutr* 91: 1348–1358
138. EFSA (2009) Scientific opinion on the substantiation of health claims related to vitamin K and maintenance of bone (ID 123, 127, 128, and 2879), blood coagulation (ID 124 and 126), and function of the heart and blood vessels (ID 124, 125 and 2880) pursuant to article 13(1) of regulation (EC) no 1924/2006. *EFSA Journal* 7: 1228–1247
139. Villa JKD, Diaz MAN, Pizzoli VR et al. (2016) Effect of vitamin K in bone metabolism and vascular calcification: a review of mechanisms of action and evidences. *Crit Rev Food Sci Nutr* [Epub ahead of print; doi: 10.1080/10408398.2016.1211616]
140. Gröber U, Reichrath J, Holick MF et al. (2014) Vitamin K: an old vitamin in a new perspective. *Dermatoendocrinol* 6: e968490
141. National Research Council. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. National Academies Press, Washington (2000)