

Einfluss von Nahrungspolyphenolen auf den Vitamin-E-Status

Studien in Modellen zunehmender Komplexität – vom physikalisch-chemischen Experiment zur Humanstudie

Jan Frank, Institut für Humanernährung und Lebensmittelkunde, Christian-Albrechts-Universität Kiel

Ernährungs-Umschau 54 (2007), S. 52 ff.

Literatur

1. Kamal-Eldin, A.; Appelqvist, L.A. The Chemistry and Antioxidant Properties of Tocopherols and Tocotrienols. *Lipids* 31:671–701; 1996.
2. Frank, J. Dietary Phenolic Compounds and Vitamin E Bioavailability – Model studies in rats and humans. Department of Food Science, Swedish University of Agricultural Sciences, Uppsala, Sweden (available online: <http://epsilon.slu.se/a446.pdf>); 2004.
3. Kayden, H.J.; Traber, M.G. Absorption, lipoprotein transport, and regulation of plasma concentrations of vitamin E in humans. *J Lipid Res* 34:343–358; 1993.
4. Parker, R.S.; Sonntag, T.J.; Swanson, J.E. Cytochrome P4503A-dependent metabolism of tocopherols and inhibition by sesamin. *Biochem Biophys Res Commun* 277:531–534; 2000.
5. Sonntag, T.J.; Parker, R.S. Cytochrome P450 omega-hydroxylase pathway of tocopherol catabolism. Novel mechanism of regulation of vitamin E status. *J Biol Chem* 277:25290–25296; 2002.
6. Burton, G.W. et al. Human plasma and tissue a-tocopherol concentrations in response to supplementation with deuterated natural and synthetic vitamin E. *Am J Clin Nutr* 67:669–684; 1998.
7. Qian, J. et al. Intracellular trafficking of vitamin E in hepatocytes: the role of tocopherol transfer protein. *J Lipid Res* 46:2072–2082; 2005.
8. Hosomi, A. et al. Affinity for alpha-tocopherol transfer protein as a determinant of the biological activities of vitamin E analogs. *FEBS Lett* 409:105–108; 1997.
9. Evans, H.M.; Bishop, K.S. On the existence of a hitherto unrecognized dietary factor essential for reproduction. *Science* 56:650–651; 1922.
10. Mason, K.E. The first two decades of vitamin E. *Fed Proc* 36:1906–1910; 1977.
11. Olcott, H.S.; Emerson, O.H. Antioxidants and the Autoxidation of Fats. IX. The Antioxidant Properties of the Tocopherols. *J Am Chem Soc* 59:1008–1009; 1937.
12. Dam, H. Vitamin E as an in vivo antioxidant. *Experientia* 1:195–213; 1952.
13. Burton, G.W.; Joyce, A.; Ingold, K.U. First proof that vitamin E is major lipid-soluble, chain-breaking antioxidant in human blood plasma. *Lancet* 320:327; 1982.
14. Rimbach, G. et al. Regulation of cell signaling by vitamin E. *Proc Nutr Soc* 61:415–425; 2002.
15. Bosco-Boinik, D. et al. Inhibition of cell proliferation by alpha-tocopherol. Role of protein kinase C. *J Biol Chem* 266:6188–6194; 1991.
16. Davies, K.J. Oxidative stress: the paradox of aerobic life. *Biochem Soc Symp* 61:1–31; 1995.
17. Keaney, J.F., Jr. et al. Obesity and systemic oxidative stress: clinical correlates of oxidative stress in the Framingham Study. *Arterioscler Thromb Vasc Biol* 23:434–439; 2003.
18. Ceriello, A.; Motz, E. Is Oxidative Stress the Pathogenic Mechanism Underlying Insulin Resistance, Diabetes, and Cardiovascular Disease? The Common Soil Hypothesis Revisited. *Arterioscler Thromb Vasc Biol*; 2004.
19. Azzi, A. Vitamin E in Cell Signaling. In: Packer, L.; Traber, M.G.; Kraemer, K.; Frei, B., eds. *The Antioxidant Vitamins C and E*. Champaign, IL, USA: AOCS Press; 2002: 195–208.
20. Saldeen, T.; Li, D.; Mehta, J. L. Differential effects of a- and g-tocopherol on low-density lipoprotein oxidation, superoxide activity, platelet aggregation and arterial thrombogenesis. *J Am Coll Cardiol* 34:1208–1215; 1999.
21. Hensley, K. et al. New Perspectives on Vitamin E: γ -Tocopherol and Carboxyethylhydroxychroman Metabolites in Biology and Medicine. *Free Radic Biol Med* 36:1–15; 2004.
22. Kontush, A. et al. Lipophilic antioxidants in blood plasma as markers of atherosclerosis: the role of a-carotene and g-tocopherol. *Atherosclerosis* 144:117–122; 1999.
23. Öhrvall, M.; Sundlöf, G.; Vessby, B. Gamma, but not alpha, tocopherol levels in serum are reduced in coronary heart disease patients. *J Intern Med* 239:111–117; 1996.
24. Kristenson, M. et al. Antioxidant state and mortality from coronary heart disease in Lithuanian and Swedish men: concomitant cross sectional study of men aged 50. *BMJ* 314:629–633; 1997.
25. Ruiz Rejón, F. et al. Plasma status of retinol, a- and g-tocopherols, and main carotenoids to first myocardial infarction: case control and follow-up study. *Nutrition* 18:26–31; 2002.
26. Gey, K.F. et al. Inverse correlation between plasma vitamin E and mortality from ischemic heart disease in cross-cultural epidemiology. *Am J Clin Nutr* 53:326S–334S; 1991.
27. Jialal, I.; Devaraj, S. Prospective Vitamin E Clinical Trials. In: Packer, L.; Traber, M. G.; Kraemer, K.; Frei, B., eds. *The Antioxidant Vitamins C and E*. Champaign, IL, USA: AOCS Press; 2002: 243–254.
28. Heinonen, O. P. et al. Prostate cancer and supplementation with a-tocopherol and b-carotene: incidence and mortality in a controlled trial. *J Natl Cancer Inst* 90:440–446; 1998.
29. Chan, J. M. et al. Supplemental vitamin E intake and prostate cancer risk in a large cohort of men in the United States. *Cancer Epidemiol Biomarkers Prev* 8:893–899; 1999.
30. Giovannucci, E. γ -Tocopherol: a new player in prostate cancer prevention? *J Natl Cancer Inst* 92:1966–1967; 2000.
31. Helzlsouer, K. J. et al. Association between a-tocopherol, g-tocopherol, selenium, and subsequent prostate cancer. *J Natl Cancer Inst* 92:2018–2023; 2000.
32. Gysin, R.; Azzi, A.; Visarius, T. g-Tocopherol inhibits human cancer cell cycle progression and cell proliferation by down-regulation of cyclins. *FASEB J* 16:1952–1954; 2002.
33. Galli, F. et al. The effect of a- and γ -tocopherol and their carboxyethyl hydroxychroman metabolites on prostate cancer cell proliferation. *Arch Biochem Biophys* 423:97–102; 2004.
34. Schwenke, D. C. Does lack of tocopherols and tocotrienols put women at increased risk of breast cancer? *J Nutr Biochem* 13:2–20; 2002.
35. Miller, E.R., 3rd et al. Meta-analysis: high-dose vitamin E supplementation may increase all-cause mortality. *Ann Intern Med* 142:37–46; 2005.
36. Lonn, E. et al. Effects of long-term vitamin E supplementation on cardiovascular events and cancer: a randomized controlled trial. *JAMA* 293:1338–1347; 2005.
37. Hatchcock, J. N. et al. Vitamins E and C are safe across a broad range of intakes. *Am J Clin Nutr* 81:736–745; 2005.
38. Scalbert, A.; Williamson, G. Dietary intake and bioavailability of polyphenols. *J Nutr* 130:2073S–2085S; 2000.
39. Leclercq, C.; Arcella, D.; Turrini, A. Estimates of the theoretical maximum daily intake of erythorbic acid, gallates, butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT) in Italy: a stepwise approach. *Food Chem Toxicol* 38:1075–1084; 2000.
40. Lampe, J. W. Health effects of vegetables and fruit: assessing mechanisms of action in human experimental studies. *Am J Clin Nutr* 70:475S–490S; 1999.
41. Segarothy, M.; Phillips, P.A. Vegetarian diet: panacea for modern lifestyle diseases? *Q J Med* 92:531–544; 1999.
42. Buettner, G. R. The pecking order of free radicals and antioxidants: lipid peroxidation, a-tocopherol, and ascorbate. *Arch Biochem Biophys* 300:535–543; 1993.
43. Nardini, M. et al. Inhibition of human low-density lipoprotein oxidation by caffeic acid and other hydroxycinnamic acid derivatives. *Free Radic Biol Med* 19:541–552; 1995.
44. Laranjinha, J.; Cadena, E. Redox cycles of caffeic acid, a-tocopherol, and ascorbate: implications for protection of low-density li-

- poproteins against oxidation. *IUBMB Life* 48:57–65; 1999.
45. *Laranjinha, J.* et al. Two related phenolic antioxidants with opposite effects on vitamin E content in low density lipoproteins oxidized by ferrylmyoglobin: consumption vs regeneration. *Arch Biochem Biophys* 323:373–381; 1995.
 46. *Laranjinha, J.A.; Almeida, L.M.; Madeira, V.M.* Reactivity of dietary phenolic acids with peroxyl radicals: antioxidant activity upon low density lipoprotein peroxidation. *Biochem Pharmacol* 48:487–494.; 1994.
 47. *Chen, J.H.; Ho, C.T.* Antioxidant activities of caffeic acid and its related hydroxycinnamic acid compounds. *J Agric Food Chem* 45:2374–2378; 1997.
 48. *Liao, K.; Yin, M.* Individual and combined antioxidant effects of seven phenolic agents in human erythrocyte membrane ghosts and phosphatidylcholine liposome systems: importance of the partition coefficient. *J Agric Food Chem* 48:2266–2270; 2000.
 49. *National Research Council Nutrient Requirements of the Laboratory Rat.* Nutrient Requirements of Laboratory Animals. Washington, D.C.: National Academic Press; 1995: 11–79.
 50. *Frank, J.* et al. Dietary (+)-Catechin and BHT Markedly Increase α -Tocopherol Concentrations in Rats by a Tocopherol- ω -Hydroxylase-Independent Mechanism. *J Nutr* 133: 3195–3199; 2003.
 51. *Pedrielli, P.; Skibsted, L. H.* Antioxidant synergy and regeneration effect of quercetin, (–)-epicatechin, and (+)-catechin on α -tocopherol in homogeneous solutions of peroxidating methyl linoleate. *J Agric Food Chem* 50:7138–7144; 2002.
 52. *Frank, J.* et al. Dietary flavonoids with a catechol structure increase α -tocopherol in rats and protect the vitamin from oxidation in vitro. *J Lipid Res* 47:2718–2725; 2006.
 53. *Pár, A.; Mézes, M.; Németh, P.; Jávor, T.* Effects of cianidanol on the blood lipid peroxide sta-
 - tus in patients with chronic hepatitis. *Int J Clin Pharmacol Res* 5:389–397; 1985.
 54. *Choi, E.J.; Chee, K.M.; Lee, B.H.* Anti- and pro-oxidant effects of chronic quercetin administration in rats. *Eur J Pharmacol* 482:281–285; 2003.
 55. *Frémont, L.* et al. Dietary flavonoids reduce lipid peroxidation in rats fed polyunsaturated or monounsaturated fat diets. *J Nutr* 128:1495–1502; 1998.
 56. *Benito, S.; Buxaderas, S.; Mitjavila, M. T.* Flavonoid metabolites and susceptibility of rat lipoproteins to oxidation. *Am J Physiol Heart Circ Physiol* 287:H2819–2824; 2004.
 57. *Rice-Evans, C.A.* et al. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radic Res* 22:375–383; 1995.
 58. *Manach, C.* et al. Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr* 81:230S–242S; 2005.
 59. *Kamal-Eldin, A.* et al. Effects of dietary phenolic compounds on tocopherol, cholesterol, and fatty acids in rats. *Lipids* 35:427–435; 2000.
 60. *Simán, C.M.; Eriksson, U.J.* Effect of butylated hydroxytoluene on α -tocopherol content in liver and adipose tissue of rats. *Toxicol Lett* 87:103–108; 1996.
 61. *Daniel, J.W.; Gage, J.C.* The absorption and excretion of butylated hydroxytoluene (BHT) in the rat. *Food Cosmet Toxicol* 3:405–415; 1965.
 62. *Rich, K.J.; Boobis, A.R.* Expression and inducibility of P450 enzymes during liver ontogeny. *Microsc Res Tech* 39:424–435; 1997.
 63. *Ikeda, S.; Tohyama, T.; Yamashita, K.* Dietary sesame seed and its lignans inhibit 2,7,8-trimethyl- 2(2'-carboxyethyl)-6-hydroxychroman excretion into urine of rats fed γ -tocopherol. *J Nutr* 132:961–966; 2002.
 64. *Cooney, R.V.* et al. Effects of dietary sesame seeds on plasma tocopherol levels. *Nutr Cancer* 39:66–71; 2001.
 65. *Lemcke-Norojärvi, M.* et al. Corn and Sesame Oils Increase Serum γ -Tocopherol Concentrations in Healthy Swedish Women. *J Nutr* 131:1195–1201; 2001.
 66. *Frank, J.; Kamal-Eldin, A.; Traber, M. G.* Consumption of Sesame Oil Muffins Decreases the Urinary Excretion of α -Tocopherol Metabolites in Humans. *Ann N Y Acad Sci* 1031:365–367; 2004.
 67. *Ross, A.B.; Aman, P.; Kamal-Eldin, A.* Identification of cereal alkylresorcinol metabolites in human urine—potential biomarkers of wholegrain wheat and rye intake. *J Chromatogr B Analyt Technol Biomed Life Sci* 809:125–130; 2004.
 68. *Ross, A.B.* et al. Cereal Alkylresorcinols Elevate α -Tocopherol Levels in Rats and Inhibit α -Tocopherol Metabolism In Vitro. *J Nutr* 134:506–510; 2004.
 69. *Ross, A.B.; Kamal-Eldin, A.; Åman, P.* Dietary alkylresorcinols: absorption, bioactivities, and possible use as biomarkers of whole-grain wheat- and rye-rich foods. *Nutr Rev* 62:81–95; 2004.
 70. *Murray, M.* Mechanisms of inhibitory and regulatory effects of methylenedioxypyhenyl compounds on cytochrome P450-dependent drug oxidation. *Curr Drug Metab* 1:67–84; 2000.
 71. *Frank, J.* et al. Dietary secoisolariciresinol diglucoside and its oligomers with 3-hydroxy-3-methyl glutaric acid decrease vitamin E levels in rats. *Br J Nutr* 92:169–176; 2004.
 72. *Frank, J.* et al. The dietary hydroxycinnamate caffeic acid and its conjugate chlorogenic acid increase vitamin E and cholesterol concentrations in Sprague-Dawley rats. *J Agric Food Chem* 51:2526–2531; 2003.
 73. *O'Byrne, D.* et al. Studies of LDL oxidation following α -, γ , or d-tocotrienyl acetate supplementation of hypercholesterolemic humans. *Free Radic Biol Med* 29:834–845; 2000.