

# Blutglukose- und -lipidsenkende Wirkung von Phytinsäure

Prof. Dr. Horst Schmandke

Ernährungs Umschau 54 (2007), S. 254 ff.

## Literatur

1. Kasper H: *Ernährungsmedizin und Diätetik*. Urban & Fischer, München, 10. Aufl. 2004
2. Lammert A et al. (2006) Kohlenhydratdrattage als einfache und effektive Therapie der Insulinresistenz. *Diabetologie* 1, S40–S41
3. Weck M et al. (1983) Ackerbohnenproteindiät bei Hypercholesterolämie. *Die Nahrung* 27, 327–333
4. Maga JA (1982) Phytate: Its chemistry, occurrence, food interactions, nutritional significance, and methods of analysis. *J Agric Food Chem* 30, 1–9
5. Reddy NR, Sathe SK, Salunkhe DH (1982) Phytates in legumes and cereals. *Adv Food Res* 28, 1–92
6. Souci SW, Fachmann W, Kraut H: *Zusammensetzung der Lebensmittel, Nährwert-Tabellen*,. Medpharm Scientific Publ, Stuttgart. 6. Aufl. 2000
7. Muschiolik G, Schmandke H: *Funktionelle Eigenschaften von Ackerbohnenprodukten (Vicia faba) – Ernährung, Biochemie und Verarbeitung*. Shaker, Aachen, 2000
8. Olendzki, BC et al. (2006) Methodology for adding glycemic index and glycemic load values to 24-hour dietary recall database. *Nutrition* 22, 1087–1095
9. Sakamoto K, Vucenik I, Shamsuddin AM (1993) [3 H] Phytic acid (inositol hexa-phosphate) is absorbed and distributed to various tissues in rats. *J Nutr* 123, 713–720
10. Nahapetian A, Young V (1980) Metabolism of <sup>14</sup>C-phytate in rats: effect of low and high dietary calcium intake. *J Nutr* 110, 1458–1472
11. Miyazawa E, Iwabuchi A, Yoshida T (1996) Phytate breakdown and apparent absorption of phosphorus, calcium and magnesium in germfree and conventionalized rats. *Nutr Res* 16, 603–616
12. Thompson LU, Button CL, Jenkins DJA (1987) Phytic acid and calcium affect the in vitro rate of navy bean starch digestion and blood glucose response in humans. *Am J Clin Nutr* 46, 467–473
13. Lee, SH et al. (2006): Dietary phytic acid lowers the blood glucose level in diabetic KK mice. *Nutr Res* 26, 474–479
14. Lee, SH (2005) Effects of dietary phytic acid on serum and hepatic lipid levels in diabetic KK mice. *Nutr Res* 25, 869–876
15. Lee HH; Lee SY; Choi YS (1999) Effects of dietary phytic acid on lipid profiles in rats of different ages. *Food Sci Biotechnol* 8, 108–112
16. Martinez-González J, Escudero I, Badimon L (2004) Simvastatin potentiates PGI<sub>2</sub> release induced by HDL in human VSMC: effect on Cox-2 up-regulation and MARK signalling pathways activated by HDL. *Atherosclerosis* 174, 305–313
17. Katayama T (1997) Effects of dietary myo-inositol or phytic acid on hepatic concentrations of lipids and hepatic activities of lipogenic enzymes in rats fed on corn starch or sucrose. *Nutr Res* 17, 721–728
18. Onomi S, Katayama T, Sato K (1999) Effects of dietary myo-inositol related compounds on sucrose-mediated hepatic lipid accumulation in rats. *Nutr Res* 19, 1401–1409
19. Okazaki Y, Setoguchi T, Katayama T (2006) Effects of dietary myo-inositol, D-chiro-inositol and L-chiro-inositol on hepatic lipids with reference to the hepatic myo-inositol status in rats fed on 1,1,1-trichloro-2,2-bis (p-chlorophenyl) ethane. *Biosci Biotechnol Biochem* 70, 2766–2770
20. Yoshida KJ et al. (2006) Genetic modification of *Bacillus subtilis* for production of D-chiro-inositol, an investigational drug candidate for treatment of type 2 diabetes and polycystic ovary syndrome. *Appl Environ Microbiol* 72, 1310–1315